



Phytochemical control of *Radopholus similis* in banana by *Azadirachta indica* and *Allium sativum* extracts

Ezra S. Bartholomew¹, Richard A.I. Brathwaite¹ & Wendy-Ann P. Isaac^{1*}

¹ Department of Food Production, The University of the West Indies, St. Augustine, Trinidad and Tobago

*email: wendy-ann.isaac@sta.uwi.edu

Abstract

The root-burrowing nematode, *Radopholus similis* is considered to be the most destructive nematode associated with banana production worldwide. This nematode can reduce plant growth and yield by more than 50% and decrease the productive life of banana fields. *R. similis* control in the Windward Islands banana industry has been based on the application of synthetic nematicides, which are now prohibited due to human health and environment hazards. One possible alternative is the utilization of plant extracts with known nematotoxic effects such as *Azadirachta indica* and *Allium sativum*. The efficacy of these phytochemicals at managing *R. similis* were assessed and compared with a synthetic nematicide, Ethoprophos (Mocap 15G) in two banana pot trials. There were 5 treatments (Control A, Control B, Neem X, Garland, and Mocap 15G)) with 6 plants per treatment. The results of the root and soil extractions showed that all treatments caused a significant reduction in *R. similis* population density ($p=0.05$), with Mocap 15G being the most effective. From the plant growth data it was observed that all treatments caused a significant increase in plant growth ($p=0.05$). No significant difference in the pseudostem length and girth, as well as, the leaf number and area were observed between treatments. The root and corm weights also showed no significant differences between treatments. The efficacy of botanical nematicides and their effect on banana production in the Windward Islands are discussed.

Keywords: Allicin, Azadirachtin, botanical control, nematostat, Trinidad and Tobago.

1. Introduction

The banana industry has been a key source of employment and foreign revenue to the Windward Islands for over eighty years (NERA, 2004). The banana trade has provided a direct living for thousands of small-scale producers and has accounted for up to 50% of the Windward Islands' total export revenue with sales of 274,000 tonnes a year to a value of US\$147m. This industry, while blessed with a warm tropical climate, is threatened by tropical storms and a complex of fungi, insects and nematodes, the latter being one of the most important limiting factors. These factors have led to a major reduction in banana production and exports to a mere 99,000 tonnes a year at a value of US\$45m (Wiltshir 2004). The root-burrowing nematode, *Radopholus similis* (Cobb1893) Thorne 1949 is

considered to be the most destructive nematode associated with banana production worldwide (Gowen et al 2005). *Radopholus similis* feeds on the cortical cells of root and corm (rhizome) tissues causing cavities to develop, which evolves as root necrosis (Brooks 2008). This in turn reduces growth and yield by more than 50%, lengthens the time to fruiting, and decreases the productive life of banana fields (Quénéhervé et al. 2006). Plant anchorage is also affected, which results in toppling or uprooting (Gowen et al 2005) Effective *R. similis* control is therefore essential for the survival of the banana industry.

Despite the fact that there has been international advocacy for organic agriculture since 1940 (Paull, 2014) the uptake of organic principles of production has been slow in Trinidad and Tobago. Although 164 countries report certified organic agriculture statistics, and some report organic banana production, no organically managed land is reported by Trinidad and Tobago (Willer & Lernoud, 2014). In the past, in Trinidad and Tobago, *Radopholus similis* control has been based mainly on the use of synthetic nematicides such as, Ethoprophos, Oxamyl and Aldicarb (Chabrier & Queneherve. 2003). These products are now prohibited due to their adverse impacts on human health and the environment (Nagaraju et al 2010). Windward Islands banana farmers are also barred from using other synthetic nematicides due to Fair-trade arrangements (Isaac et al., 2007). Plant-based pest control agents have long been touted as alternatives to synthetic nematicides (Javed et al 2006). Such phytochemicals reputedly pose little threat to the environment or to human health and their use is approved under organic and Fair-trade arrangements (Amadioha, 2003). However, the adoption of such a management strategy has been met with scepticism by banana producers who question the efficacy and consistency of these phytochemicals (Villanueva 2005). In this study, phytochemicals derived from *Azadirachta indica* and *Allium sativum* extracts were assessed and compared with Ethoprophos for their effectiveness at reducing *R. similis* population density in the roots and soil, and preventing banana growth and yield losses.

2. Materials and Methods

Two pot trials were conducted throughout 2009 and 2010 at the University of the West Indies Field Station (UFS) situated in Valsayn, Trinidad (10° 39' 0" N/ 61° 25' 0" W). The average monthly temperature and precipitation at UFS were 27.2°C and 1719.5mm, respectively for the length of the study. The growing medium comprised of sterilized Fluventic Eutropept soil (River Estate Loam), with a CEC of 4.8 cmol/kg and a pH of 6.5.

2.1 Experimental Design

Each trial consisted of 30 banana plants (cv. Lacatan), planted in 50 L plastic-drums and spaced 2×2m apart. The drums were arranged in a completely randomized design with five treatments (Control A, Control B, Neem X, Garland, and Mocap 15G) and six replicates. Plants were grown for 1 month and then inoculated with 500 *R. similis* following the procedures of Speijer & De Waele (1997). One month after inoculation, nematicidal treatments were applied as a soil drench (1L) at the base of the pseudostem. The active ingredients, sources, application rates and frequencies of each treatment are given in Table 1. Agronomical management was done according to the protocols recommended in the Windward Islands for banana production, which included fertilization, irrigation, pruning, propping, de-suckering, and pest and disease management (Paul et al 1993).

Table 1: Active Ingredients, sources, application rates and frequencies of treatments

Treatments (Trade Names)	Active Ingredients (Common Names)	Parent company	Application rate	Application frequency
Control A	No <i>R. similis</i>	-	-	-
Control B	<i>R. similis</i> only	-	-	-
Neem X®	Azadirachtin (3000 ppm)	Marketing Arm International	2.7ml/L	7 days intervals for 3 applications
Garland®	Allicin (<1 ppm)	OMEX Agriculture Ltd	2.16ml/L	7 days intervals for 3 applications
Mocap 15G®	Ethoprophos	Bayer Crop Science	15g/m ²	7 days intervals for 3 applications

2.2 Data Collection

i. *Radopholus similis* Population Density

An estimation of the *R. similis* population density was done using extraction methodologies described by Southey (1986) for the roots, and Whitehead & Hemming (1965) for the soil. The blender nematode filter extraction method was used to extract nematodes from 15g of roots while the Whitehead tray method was modified to extract nematodes from 200ml of soil. The collected nematode sample was identified and counted in three 1ml aliquots out of a 10ml aqueous suspension using a stereoscopic microscope. All vermiform stages (juveniles and adults) were counted.

ii. Root Necrosis Index and Banana Root and Corm Fresh Weight

At the end of each trial the plants were excavated, their roots and corms cleaned to remove soil particles and their root necrosis index (RNI) and root and corm fresh weights (RCFW) determined. A modified Bridge & Gowen (1993) root necrosis index was used to determine root rotting on a 0 to 4 scale [0 = no damage; 1 = <25% of total root cortex with necrosis; 2 = 26–50% of the total root cortex with necrosis; 3 = 51–75% of total root cortex with necrosis; 4 = >75% of total root cortex with necrosis]. Fresh weight (kg) was determined using a Rebure pocket balance (Germany).

iii. Plant Growth

Throughout the study the following crop measurements were recorded weekly:

- The pseudostem length (cm) measured from the point of the lowest leaf to the base of the pseudostem.
- The pseudostem girth/circumference (cm) measured from a point at half the pseudostem length.
- The total number of fully opened functional leaves (Fogain 2000).
- Leaf area (LA) predicted with the regression models: $LA = 0.0266 + (L \times W \times 0.7629)$ ($r = 0.98$), where L = leaf length and W = leaf width. The third leaf from the top of the plants was selected as the standard leaf for measurement (Potder & Pauer 1990).

2.3 Data Analysis

Differences in nematode density, necrosis index, roots and corm fresh weight, and growth parameters were analyzed with the use of analysis of variance (ANOVA). Pearson

correlation was used to determine the strength of relationship between the *R. similis* density in the roots and the necrosis index. Prior to analyses, variables were tested for homogeneity of variances and normality, and data found to be non-homogenous were either Log10 ($X + 1$) or square root transformed before statistical analysis. Non-transformed means were reported in Figures and Tables and only significant differences ($P \leq 0.05$) will be discussed unless stated otherwise. All statistical analysis was performed using the statistical software Minitab® 16.1.1 (Minitab Inc).

3. Results and Discussion

3.1 *Radopholus similis* Population Density

All nematicides were successful at reducing *R. similis* population density in the root and soil. Plants treated with Ethoprophos had a lower *R. similis* density than those treated with *Azadirachta indica* and *A. sativum* extracts (Fig.1). Several studies have reported on the notable effects of Ethoprophos in *Musa* spp. which acts as a nematostat in low concentrations (Stirling & Pattison 2008; Quénéhervé, 2009; Radwan et al 2012). However, to maintain its high effectiveness, frequent applications were required, which increases its negative effects on the environment and human health (Sipes & Schmitt 1995).

Plants treated with *Azadirachta indica* had a high *R. similis* density in the soil (570) but a lower density in roots (275). This disparity was due the anti-feedant properties of Azadirachtin, the phytochemical found in *Azadirachta indica* (Sidhu 2003). Rehman et al (2009) suggested that Azadirachtin can induce nematostasis, a process that inhibits nematodes from invading plants without directly killing them. The phytochemicals found in *Allium sativum* (Allicin) demonstrated a similar but less effective anti-feedant effect and was inconsistent at reducing *R. similis* density in the roots. This inconsistency may be due to the life cycle of *R. similis*, which can be completed in the root, without any stage in the soil. Thus preventing exposure to the Allicin applied in the soil (Araya 2003).




Figure 1. Effects of nematicidal treatments on *R. similis* density in the roots and soil of banana plants. Values are the average of 6 replicates. Bars with the same letter are not significantly different ($P > 0.05$).

Endo-parasitic nematodes, such as *R. similis* were expected to be more prevalent in the roots than the soil. This study showed a deviating trend which can be attributed to the nematostatic activities of the treatments along with the inefficiencies of Southey's (1986) extraction technique at recovering *R. similis* from the roots. This is a major deficiency in Southey's technique since endo-parasitic nematodes were unlikely to migrate from healthy/necrosis free root tissues to be extracted by this technique. Therefore, the estimated nematode density in the roots may have been lower than the actual nematode density.

3.2 Root Necrosis Index

The differences in the RNI between treatments were statistically significant ($P < 0.05$) in both trials, with Control A having the lowest index value and Control B the highest (Table 2). The Pearson correlation between *R. similis* density and the root necrosis were positive in both trials (Trial 1: $r^2=0.574$; Trial 2: $r^2=0.190$). Therefore, increases in the *R. similis* density in the roots will result in an increase in the root necrosis. The strength of the relationship was however inconsistent and the weak correlation in Trial 2 supports the assumption that highly necrotized root tissue may have a lower *R. similis* density. Dosselaere (2003) indicated that nematodes may move out of resource scarce necrotic banana root tissue and reinvaded healthy tissue. Plants treated with *Azadirachta indica* had the lowest R.N.I. in both trials due to the nematostatic activities of Azadirachtin, which inhibits nematodes invasion into the roots. *Allium sativum* had similar but less effective nematostatic activities, while the efficacy of Ethoprophos was inconsistent.

Table 2: Mean roots necrosis index values and root and corm fresh weight (kg).

Treatments	Mean Necrosis Index		Mean Root and Corm Weight kg)	
	Trial 1 (SD)	Trial 2 (SD)	Trial 1 (SD)	Trial 2 (SD)
Control A	1.33 (0.52)	1.83 (0.75)	2.2 a (0.75)	2.1 a (1.11)
Control B	2.67 (0.75)	2.17 (0.75)	1.4 b (0.58)	1.6 b (0.80)
Garland	2.17 (0.82)	1.83 (0.98)	1.6 b (0.60)	2.2 a (1.24)
Neem-X	1.83 (0.75)	1.67 (0.82)	1.8 ab (0.71)	2.1 a (0.56)
Mocap	1.83 (0.98)	2.17 (0.98)	2.0 a (0.33)	2.1 a (1.14)

3.3 Banana Root and Corm Fresh Weight

All treatments were effective at maintaining healthy root and corm. Ethoprophos performed consistently in both trials and recorded the highest RCFW among the treatments. The anti-feedant effect of Azadirachtin and Allicin were also consistent and comparable to Ethoprophos. *Radopholus similis* population densities in the roots and corm tissues had a direct effect on its RCFW. Moens et al (2003) found a linear reduction of root weight when *R. similis* was inoculated at increasing densities. Marin et al. (1999) and Sarah (2000) also found decreases in root weight, ranging from 8-80%, several weeks after inoculation with *R. similis*. The data obtained in this study were similar to those reported in the literature, as Control A had the highest sum of RCFW while Control B had the lowest (Table 2). Therefore, RCFW shows some potential for used as a rapid technique for estimating *R. similis* density (Moens et al 2003). However, depending solely

on RCFW to determine the implementation of nematode control strategies may not be advisable, as root and corm weight may vary due to factors other than nematode density, such as the banana weevil borer (*Cosmopolites sordidus*) and plant nutrient (Sarah 2000).

3.4 Plant Growth

The banana growth data confirmed that the phytochemical treatments were comparable to Ethoprophos at preventing plant growth losses, as no significant differences were observed in both trials (Table 3). This may also be due to the ability of banana cv. Lacatan to withstand *R. similis* infestation. Therefore, plant sensitivity or tolerance must be considered since nematode tolerance has been identified in the Musa gene pool (Dochez et al 2006) The selection of nematode tolerant banana cultivars can play a very important role in nematode management and is ranked as one of the ideotype requirements for the acceptance of banana hybrids (Tenkauan & Swennen 2004). On the other hand, a tolerant but susceptible banana plant is of limited value, as nematode reproduction may increase the population density beyond the damage threshold (Cook & Starr 2006, 372). Therefore, in the absence of true resistance, incorporating tolerant plants with phytochemicals may be an effective strategy for preventing the *R. similis* density from crossing the plant damage threshold and causing yield losses.

Table 3: Effects of the treatments on the growth parameters of banana plant

Treatments	Pseudostem		Leaf	
	Length (cm)	Girth (cm)	Numbers	Area (cm ²)
Control A	95(0.52)	28 (0.75)	6 (0.75)	2673 (1.11)
Control B	77 (0.75)	23.6 (0.75)	7.5 (0.58)	2182 (0.80)
Garland	81.8 (0.82)	26.2 (0.98)	6.7 (0.60)	2363 (1.24)
Neem-X	84.8 (0.75)	27.5 (0.82)	6.7 (0.71)	2498 (0.56)
Mocap	92 (0.98)	28.5 (0.98)	6.5 (0.33)	2691 (1.14)

4. Conclusion

The application of phytochemicals (Azadirachtin and Allicin) as alternatives to synthetic nematicides was effective and comparable to Ethoprophos at preventing plant growth and yield losses. This will be beneficial to banana producers by decreasing the time for fruiting and increasing the productive life of the fields. Plants treated with Azadirachtin also had a low root necrosis index which may result in fewer plants toppling and uprooting. However, the phytochemicals were less effective at reducing *R. similis* density in the soil which leaves the plants vulnerable to future infestations unless other nematode management strategies are adopted.

In the pursuit of more benign pest management and more consumer-acceptable solutions in banana production it would be appropriate for more research to be directed to organic research and some attention paid to the lessons learned in other countries which are pursuing the expansion of their own organic banana production.

Acknowledgement

The authors thank the manager and staff of the UFS, for their kind cooperation and support in executing these trials. Thanks to the academic and technical staff of the Department of Food Production, the University of the West Indies, St. Augustine, Trinidad and Tobago for assistance.

References

- Araya, M. 2003. Current status of management of nematodes in banana (*Musa AAA*) and plantain (*Musa AAB*) in the American tropics. In: Rivas, G. & Rosales, F. (eds.) *Conventional and Alternative Management of Black Sigatoka, Nematodes and Other Pests Associated to Musa Cultivation in the Tropics*. International Plant Genetics Resources Institute, Ecuador. pp. 79-102.
- Amadioha, A. C. 2003. Evaluation of some plant leaf extracts against *Colletotrichum lindemuthianum* in cowpea. *Actaphyto pathologica et Entomologica Hungarica*, 38(3-4): 259-265.
- Brooks, F.E. 2008. Burrowing Nematode. The Plant Health Instructor. <http://www.apsnet.org/edcenter/intropp/lessons/Nematodes/Pages/Burrowingnematode.aspx> (Viewed: 29 August, 2014)
- Bridge, J. & Gowen, S.R. 1993. Visual assessment of plant parasitic nematode and weevil damage on bananas and plantain." In: Gold, C. S. & Gemmill, B (eds.) *Biological and Integrated Control of Highland Banana and Plantain Pests and Diseases*. Research Coordination Meeting, 12-14 November 1991. Cotonou, Benin. IITA, Nigeria. pp. 147-154.
- Chabrier, C. & Queneherve, P. 2003. Control of the burrowing nematode (*Radopholus similis* Cobb) on banana: impact of the banana field destruction method on the efficiency of the following fallow. *Crop Protection*, 22: 121–127
- Cook, R. & Starr, J. L. 2006. Resistant Cultivars. In: Perry, R. N. & Moens, M. (eds). *Plant Nematology*, CAB International, Wallingford, UK. pp. 370-391.
- Dochez, C., Tenkouano, A., Ortiz, R., Whyte, J.B.A. & De Waele, D. 2006. New sources of resistance to *Radopholus similis* in *Musa* germplasm from Asia. *Australasian J Plant Path*, 35: 481 – 485.
- Dosselaere, N., Araya, M. & De Waele, D. 2003. Effect of pot volume on root growth, *Radopholus similis* reproductive potential and its damage on bananas. *INFOMUSA 12* (1):
- Fogain, R. 2000. Effect of *Radopholus similis* on plant growth and yield of plantains (*Musa*, AAB). *Nematology 2*(2): 129-133.
- Gowen, S. R., Quénéhervé, P. & Fogain, R. 2005. Nematode parasites of bananas and plantains. In: Luc, M., Sikora, R.A. & Bridge J. (eds.) *Plant parasitic nematodes in subtropical and tropical agriculture*, 2nd edition. CAB International Edit., Wallingford, UK. pp. 611–643.
- Isaac, W.A.P., Brathwaite, R.A.I., Ganpat, W.P., & Bekele, I. 2007. The impact of selected cover crops on soil fertility, weed and nematode suppression through Farmer Participatory Research by FairTrade banana growers in St. Vincent and the Grenadines. *World Journal of Agricultural Sciences 3* (1): 1-9
- Javed, N., Gowen, S., Inam-ul-Haq, R M., Abdullah, K. & Shahina, F. 2006. Systemic and persistent effect of neem (*Azadirachta indica*) formulations against root-knot nematodes, *Meloidogyne javanica* and their storage life. *Crop Protection*, 26(7): 911-916.
- Marin, D. H., Barker, K. R., Kaplan, D. T., Sutton, T. B. & Opperman, C. H. 1999. Aggressiveness and damage potential of Central American and Caribbean populations of *Radopholus* spp. in banana, *Journal of Nematology 31*: 241-247
- Moens, T., Araya, M. & De Waele, D. 2003. Correlations between nematode numbers and damage to banana (*Musa AAA*) roots under commercial conditions. *Nematropica*, 31:55-65.
- Nagaraju, M., Karemegam, N. & Kadalmani, B. 2010. Eco-Friendly Management of Root-knot nematode *Meloidogyne incognita* using Organic amendments on Tomato. *International Journal of Research in Pharmaceutical Sciences*, 1 (4): 530-532.

- NERA (National Economic Research Associates) and Oxford Policy Management. 2004. Addressing the Impact of Preference Erosion in Bananas on Caribbean Countries. London: DFID.
- Paul, N.C., Glasgow, G. & Blanchard, M. (eds.) 1993. Banana growers' manual: A guide to successful banana production in the Windward Islands, 4th ed. Voice Publishing Company: St. Lucia, Windward Islands. International Network for the Improvement of Banana and Plantain, Montpellier, France. pp. 32-44.
- Paull, J. 2014. Lord Northbourne, the man who invented organic farming, a biography. *Journal of Organic Systems*, 9(1), 31-53.
- Quénéhervé, P., Chabrier, C., Auwerkerken, A., Topart, P., Martiny, B. & Martie-Luce, S. 2006. Status of weeds as reservoirs of plant parasitic nematodes in banana fields in Martinique. *Crop Protection*, 25:860–867.
- Quénéhervé, P. 2009. Integrated management of banana nematodes. In: Ciancio, A. & Mukerji, K.G. (eds.) *Integrated management of fruit crops and forest nematodes*. Springer, Dordrecht, The Netherlands. pp. 3-61
- Radwan, M.A., Farrag, S.A.A., Abu-Elamayem, M.M. & Ahmed, N.S. 2012. Efficacy of some granular nematicides against root-knot nematode, *Meloidogyne incognita* associated with tomato. *Pakistan Journal of Nematology*, 30 (1): 41-47
- Rehma, A.U., Javed, N., Ahmad, R. & Shahid, M. 2009. Protective and Curative effect of Bio-products against the invasion and development of Root-knot nematode in Tomato. *Pakistan Journal of Phytopathology*, 21(1): 37-40
- Sarah, J. L. 2000. Burrowing nematode. In: Jones, D. R (ed.) *Diseases of banana, abacá and enset*. CAB International, Wallingford, UK. pp. 295-303.
- Sidhu, O. P., Kumar, V. & Behl, H. M. 2003. Variability in Neem (*Azadirachta indica*) with respect to azadirachtin content. *Journal of Agricultural and Food Chemistry*, 51(4):910-5.
- Sipes, B., & Schmitt, D. P. 1995. Evaluation of Ethoprop and Tetrathiocarbonate for Reniform Nematode Control in Pineapple. *Supplement to the Journal of Nematology*, 27(4S):639-644.
- Southey, J.F. 1986. Laboratory methods for work with plant and soil nematodes. Reference book 402, 6th edition. Ministry of Agriculture, Fisheries and Food, Her Majesty's Stationery Office, London. pp.202
- Speijer, P.R. & De Waele, D. 1997. Screening of *Musa* germplasm for resistance and tolerance to nematodes, INIBAP Technical Guidelines 1, International Network for the Improvement of Banana and Plantain, Montpellier, France. pp. 47
- Stirling, G.R. & Pattison, A.B. 2008. Beyond chemical dependency for managing plant-parasitic nematodes: Examples from the banana, pineapple and vegetable industries of tropical and subtropical Australia. *Australasian Plant Pathology*, 37 (3): 254-267.
- Tenkauano, A. & Swennen, R.L. 2004. Progress in breeding and delivering improved plantain and banana to African farmers. *Chronica Horticulturae*, 44(1): 9-15.
- Villanueva, L.M. 2005. Status of nematode problem affecting banana in the Philippines. In: Dela Cruz, F.S., Van den Bergh, I., Waele, D., Hautea, D.M. & Molina, A.B. (eds.) *Towards management of Musa nematodes in Asia and the Pacific: Country reports*. INIBAP, Los Baños, Laguna (PHL). pp.51-59
- Whitehead, A.G. & Hemming, J. R. 1965. A comparison of some quantitative methods of extracting small vermiform nematodes from soil. *Annals of Applied Biology*, 55:25-38.
- Willer, H., & Lernoud, J. (Eds.). 2014. *The World of Organic Agriculture: Statistics and Emerging Trends 2014*: Frick, Switzerland: Research Institute of Organic Agriculture (FiBL) & Bonn: International Federation of Organic Agriculture Movements (IFOAM).

Wiltshir, A. 2004. Fairtrade Bananas Impact Study. Dominica, Windward Islands. http://www.fairtrade.net/fileadmin/user_upload/content/2009/resources/2004_Fairtrade_Bananas_Impact_Study.pdf (Viewed: 29 August, 2014).

This is an open access article distributed under the terms of the Creative Commons Attribution License which permits unrestricted use, distribution, and reproduction in any medium, provided that the author/s and the Journal of Organics are credited.